

Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More...

International Food Research Journal  
Volume 25, Issue 6, 2018, Pages 2307-2312

Screening of lutein content in several fresh-water microalgae (Article)

Sallehudin, N.J.<sup>a</sup>, Raus, R.A.<sup>b</sup>, Mustapa, M.<sup>a</sup>, Othman, R.<sup>b</sup>, Mel, M.<sup>a</sup>

<sup>a</sup>Department of Biotechnology Engineering, Kulliyyah of Engineering, International Islamic University Malaysia, Kuala Lumpur, 50728, Malaysia  
<sup>b</sup>Department of Landscape Architecture, Kulliyyah of Architecture and Environmental Design, International Islamic University Malaysia, Kuala Lumpur, 50728, Malaysia

Abstract

View references (23)

Lutein is the most abundant xanthophylls in the photosynthetic organisms and its dietary intake is thought to protect against the development of age-related macular degeneration. Microalgae are predicted to be the major contributors of lutein in the future and to date, various species of microalgae has been analyzed for its lutein content. This study examined the lutein content of six chlorophycean microalgae species grown in the laboratory culture under the comparable environmental condition in which the population doubling time ( $t_d$ ) as well as the maximum growth rate ( $\mu_{max}$ ) for each species was evaluated. Using batch culture operation grown under the heteroautotrophic condition, *Scenedesmus dimorphus* was observed to accumulate the highest lutein content (60.11 mg/g) with maximal biomass of 0.349 g/L culture. Based on the observations available, it appears that strains of unicellular microalgae of smaller size have appreciably higher intrinsic growth rates than algae of the larger size (*Chlorella* sp. B > *Chlorella* sp. TLL > *Scenedesmus dimorphus* > *Scenedesmus obliquus* TLL > *Desmodesmus* sp. TH > *Ankistrodesmus* sp. TH). © All Rights Reserved.

SciVal Topic Prominence

Topic: astaxanthin | Xanthophylls | astaxanthin accumulation  
Prominence percentile: 97.798

Reaxys Database Information

View Compounds

Author keywords

Ankistrodesmus Chlorella Chlorophycean microalgae Desmodesmus Freshwater microalgae Growth rate Lutein Scenedesmus

Funding details

Funding sponsor	Funding number	Acronym
International Islamic University Malaysia		IIUM
Universiti Teknologi MARA		UiTM
Faculty of Pharmacy, Cairo University		

Metrics

0 Citations in Scopus  
0 Field-Weighted Citation Impact

PlumX Metrics  
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:  
Set citation alert >  
Set citation feed >

Related documents

Effects of nitrogen source availability and bioreactor operating strategies on lutein production with *Scenedesmus obliquus* FSP-3  
Ho, S.-H. , Xie, Y. , Chan, M.-C. (2015) *Bioresource Technology*  
Lutein production with wild-type and mutant strains of *Chlorella sorokiniana* MB-1 under mixotrophic growth  
Chen, J.-H. , Chen, C.-Y. , Chang, J.-S. (2017) *Journal of the Taiwan Institute of Chemical Engineers*  
Spectral characterization of  $\beta$ ,  $\epsilon$ -carotene-3, 3'-diol (lutein) from marine microalgae *Chlorella salina*  
Gayathri, S. , Rajasree, S.R.R. , Kirubakaran, R. (2016) *Renewable Energy*  
View all related documents based on references  
Find more related documents in Scopus based on:

Ministry of Higher Education, Malaysia

MOHE

## Funding text

0011 under the Ministry of Higher Education, Malaysia and by the Biotechnology Engineering Department, Kuliyah of Engineering, International Islamic University Malaysia, Gombak. The utility for freeze dryer machine was supported by Faculty of Pharmacy, University Teknologi MARA (UiTM)

ISSN: 19854668

Source Type: Journal



Original language: English

Document Type: Article

Publisher: Universiti Putra Malaysia

## References (23)

View in search results format &gt;

☐ All   [Export](#)    Print    E-mail   [Save to PDF](#)   [Create bibliography](#)

- ☐ 1 Agusti, S., Duarte, C.M., Kalff, J.  
Algal cell size and the maximum density and biomass of phytoplankton (Open Access)

(1987) *Limnology and Oceanography*, 32 (4), pp. 983-986. Cited 39 times.  
doi: 10.4319/lo.1987.32.4.0983

[View at Publisher](#)

- ☐ 2 Breithaupt, D.E., Wirt, U., Bamedi, A.  
Differentiation between lutein monoester regioisomers and detection of lutein diesters from marigold flowers (*Tagetes erecta* L.) and Several fruits by liquid chromatography-mass spectrometry

(2002) *Journal of Agricultural and Food Chemistry*, 50 (1), pp. 66-70. Cited 116 times.  
doi: 10.1021/jf010970l

[View at Publisher](#)

- ☐ 3 Cerón, M.C., Campos, I., Sánchez, J.F., Acien, F.G., Molina, E., Fernández-Sevilla, J.M.  
Recovery of lutein from microalgae biomass: Development of a process for *Scenedesmus almeriensis* biomass

(2008) *Journal of Agricultural and Food Chemistry*, 56 (24), pp. 11761-11766. Cited 72 times.  
<http://pubs.acs.org/doi/pdfplus/10.1021/jf8025875>  
doi: 10.1021/jf8025875

[View at Publisher](#)

- ☐ 4 Chan, M.-C., Ho, S.-H., Lee, D.-J., Chen, C.-Y., Huang, C.-C., Chang, J.-S.  
Characterization, extraction and purification of lutein produced by an indigenous microalga *Scenedesmus obliquus* CNW-N

(2013) *Biochemical Engineering Journal*, 78, pp. 24-31. Cited 40 times.  
doi: 10.1016/j.bej.2012.11.017

[View at Publisher](#)